

Stochastic Process Modeling of an Infectious Disease in a Finite Population

Hideaki Takagi, Professor Emeritus, University of Tsukuba, Japan
Brian L. Mark, George Mason University, Fairfax, Virginia, U.S.A.

Abstract

A birth-and-death process model with a finite state space is proposed and studied for an infectious disease spreading with several solution techniques such as a planar partial differential equation solved by Lagrange's method, direct inversion of Laplace transforms of the forward Kolmogorov equations, Syski's spectral decomposition for general continuous-time Markov processes, and Karlin-McGregor's symmetrization of birth-and-death processes along with simple numerical examples.

Table of Contents

1. Introduction
2. Kolmogorov Equations for the Birth-and-Death Process
3. Numerical Solution by Mathematica DSolve Function
4. Special Case of No Internal Infection
5. Inversion of Laplace Transforms
6. Syski's Spectral Decomposition
7. Karlin-McGregor's Symmetrization
8. References and Acknowledgments

1 Introduction

So-called SIR models have been used as major mathematical tools for quantitative analysis and prediction of the recent world-wide spread of COVID-19 without much success. The SIR-type models are based on a set of simultaneous nonlinear ordinary differential equations in several non-probabilistic variables. In principle, it would be very difficult for theoretically deterministic approaches to account for the large statistical variation in the number of infected people observed in different cities and countries with similar conditions around the world. The deterministic models could, at most, only yield average values and explain the time-varying characteristics of the pandemic in the past.

The spreading of an infectious disease is a probabilistic phenomenon because the contact with infectious persons does not necessarily cause the infection. It cannot be either certain that the vaccination suppresses infection with 100 percent of success. Therefore, we should not depend on deterministic models for important prediction.

Stochastic process models have been exploited for population biology and epidemics mostly assuming an infinite population. In the present paper, we focus on the analysis of a finite population model. A real case of the infection in a finite population is the COVID-19 outbreak among passengers and crew on Diamond Princess cruise ship in Yokohama Port, Japan, in early 2020 [7]. We believe that exploration of stochastic techniques for epidemic models will contribute to academia as well as to the society in general with significant impact.

2 Kolmogorov Equations for the Birth-and-Death Process

Our basic model for the spreading of an infectious disease is a continuous-time birth-and-death process of a stochastic variable $I(t)$, the number of infected persons at time t (≥ 0) in a discrete state space between 0 and N inclusive. We assume the three positive, constant parameters as follows:

- λ (birth rate) : Mean number of persons who get infected from others per unit time,
- μ (removal rate) : Mean number of persons who die or recover per unit time,
- ν (immigration rate) : Mean number of persons who become infectious per unit time.

We study the probability $P_n(t) \triangleq P\{I(t) = n \mid P(0) = I_0\}$, $0 \leq n \leq N$. For a short interval $\Delta t \ll t$, we consider the events occurring during an infinitesimal interval $(t, t + \Delta t]$. We get the forward Chapman-Kolmogorov equations for the process $\{I(t); t \geq 0\}$ as

$$P_0(t + \Delta t) = (1 - N\nu\Delta t)P_0(t) + \mu\Delta tP_1(t) + o(\Delta t), \quad (1)$$

$$P_n(t + \Delta t) = \{1 - [n(\lambda + \mu) + (N - n)\nu]\Delta t\}P_n(t) + (n + 1)\mu\Delta tP_{n+1}(t) + \{(n - 1)\lambda + (N - n + 1)\nu\}\Delta tP_{n-1}(t) + o(\Delta t) \quad 1 \leq n \leq N - 1, \quad (2)$$

$$P_N(t + \Delta t) = (1 - N\mu\Delta t)P_N(t) + \{(N - 1)\lambda + \nu\}\Delta tP_{N-1}(t) + o(\Delta t). \quad (3)$$

By moving some terms, dividing both sides by Δt , and making $\Delta t \rightarrow 0$, we obtain the following set of simultaneous differential-difference equations for $\{P_n(t); 0 \leq n \leq N\}$:

$$\frac{dP_0(t)}{dt} = -N\nu P_0(t) + \mu P_1(t), \quad (4)$$

$$\begin{aligned} \frac{dP_n(t)}{dt} &= -\{n(\lambda + \mu) + (N - n)\nu\}P_n(t) + (n + 1)\mu P_{n+1}(t) \\ &+ \{(n - 1)\lambda + (N - n + 1)\nu\}P_{n-1}(t) \quad 1 \leq n \leq N - 1, \end{aligned} \quad (5)$$

$$\frac{dP_N(t)}{dt} = -N\mu P_N(t) + \{(N - 1)\lambda + \nu\}P_{N-1}(t). \quad (6)$$

In addition, we impose the normalization condition: $\sum_{n=0}^N P_n(t) = 1$.

Let us introduce the Probability Generating Function (PGF) for $I(t)$ by

$$G(z, t) \triangleq E \left[z^{I(t)} \right] = \sum_{n=0}^N P_n(t) z^n \quad ; \quad \frac{\partial G(z, t)}{\partial z} = \sum_{n=1}^N n P_n(t) z^{n-1}. \quad (7)$$

Then, we get the following planar partial differential equation:

$$\frac{\partial G(z, t)}{\partial t} - (z - 1) \{ (\lambda - \nu) z - \mu \} \frac{\partial G(z, t)}{\partial z} = N \nu (z - 1) G(z, t) + N \lambda P_N(t) z^N. \quad (8)$$

So far, we have been unable to solve this equation analytically due to the last term on the right-hand side. Therefore, in the next section, we show the numerical solution using the “DSolve” function of Mathematica.

3 Numerical Solution by Mathematica DSolve Function

In Table 1, we tabulate the probabilities $\{P_n(t); 0 \leq n \leq 8\}$, the mean $E[I(t)]$, and the variance $\text{Var}[I(t)]$ by assuming $\lambda = 0.5$. The numerical values are obtained by the solution of the set of simultaneous ordinary differential equations given in eqs. (4)–(6) by Mathematica “DSolve” function.

Table 1: Probabilities $\{P_n(t); 0 \leq n \leq 8\}$, the mean $E[I(t)]$, and the variance $\text{Var}[I(t)]$. Parameters: $N = 8, I(0) = 5, \lambda = 0.5, \nu = 0.2, \mu = 0.1$.

t	$P_0(t)$	$P_1(t)$	$P_2(t)$	$P_3(t)$	$P_4(t)$	$P_5(t)$	$P_6(t)$	$P_7(t)$	$P_8(t)$	$E[I(t)]$	$\text{Var}[I(t)]$
0	0	0	0	0	0	1.000000	0	0	0	5.00000	0
1	≈ 0	0.000037	0.000646	0.006009	0.031521	0.093509	0.161557	0.225604	0.481116	7.01049	1.413950
2	≈ 0	0.000059	0.000514	0.002631	0.009265	0.024756	0.061208	0.181213	0.718915	7.56685	0.698757
3	≈ 0	0.000038	0.000239	0.001004	0.003402	0.011135	0.040970	0.172656	0.770553	7.69165	0.431854
4	≈ 0	0.000019	0.000105	0.000450	0.001831	0.007923	0.036509	0.170672	0.782489	7.72219	0.356205
5	≈ 0	0.000009	0.000054	0.000274	0.001392	0.007097	0.035420	0.170236	0.785516	7.73030	0.334643
6	≈ 0	0.000001	0.000036	0.000219	0.001265	0.006871	0.035133	0.170130	0.786340	7.73258	0.328290
7	≈ 0	0.000003	0.000030	0.000202	0.001227	0.006806	0.035053	0.107103	0.786576	7.73325	0.326360
8	≈ 0	0.000003	0.000027	0.000196	0.001216	0.006787	0.035029	0.170095	0.786646	7.73345	0.325759
9	≈ 0	0.000003	0.000027	0.000195	0.001212	0.006781	0.035022	0.170093	0.786667	7.73352	0.325568
10	≈ 0	0.000003	0.000027	0.000193	0.001211	0.006779	0.035020	0.170093	0.786674	7.73354	0.325507

4 Special Case of No Internal Infection

In this section, we consider a special case $\lambda = 0$, which means that the internal infection never occurs by means of perfect management of infected persons. In such a case, we

are to solve the partial differential equation ¹

$$\frac{\partial G(z, t)}{\partial t} + (z - 1)(\nu z + \mu) \frac{\partial G(z, t)}{\partial z} = N\nu(z - 1)G(z, t). \quad (9)$$

Then we can use well-known Lagrange's method. The corresponding set of auxiliary differential equations defining the normal at a point on the solution surface is given by

$$\frac{dt}{1} = \frac{dz}{(z - 1)(\nu z + \mu)} = \frac{dG(z, t)}{N\nu(z - 1)G(z, t)}. \quad (10)$$

Following a standard procedure, we obtain the PGF for $I(t)$ as follows:

$$G(z, t) = [q(t)z + 1 - q(t)]^{I_0} [r(t)z + 1 - r(t)]^{N-I_0}, \quad (11)$$

which is a polynomial in z of degree N , where

$$q(t) \triangleq \frac{\nu + \mu e^{-(\nu+\mu)t}}{\nu + \mu} \quad ; \quad r(t) \triangleq \frac{\nu (1 - e^{-(\nu+\mu)t})}{\nu + \mu}. \quad (12)$$

Therefore, the PGF $G(z, t)$ is the product of the PGF's for two statistically independent processes, each being the binomial (or Bernoulli) distribution.

4.1 Mean and Variance

Since the two binomial distributions are independent, the mean of $I(t)$ is given by the sum of the means for each distribution. Similarly, the variance of $I(t)$ is given by the sum of the variances for each distribution.

$$E[I(t)] = I_0 e^{-(\nu+\mu)t} + \frac{N\nu}{\nu + \mu} (1 - e^{-(\nu+\mu)t}), \quad (13)$$

$$\text{Var}[I(t)] = I_0 \frac{\nu - \mu}{\nu + \mu} e^{-(\nu+\mu)t} (1 - e^{-(\nu+\mu)t}) + \frac{N\nu^2}{(\nu + \mu)^2} (1 - e^{-2(\mu+\nu)t}). \quad (14)$$

4.2 Probability Mass Function

The probability mass function (PMF) of $I(t)$ is given by the convolution of two binomial distributions (Bailey [1, p.64], Kobayashi and Ren [5], Syski [6, p.138]).

$$P_0(t) = [1 - q(t)]^{I_0} [1 - r(t)]^{N-I_0} = \left[\frac{\mu (1 - e^{-(\nu+\mu)t})}{\nu + \mu} \right]^{I_0} \left[\frac{\mu + \nu e^{-(\nu+\mu)t}}{\nu + \mu} \right]^{N-I_0}, \quad (15)$$

¹Solution to the equivalent partial differential equation is shown in [5] and [6, p.138].

$$P_n(t) = \sum_{i=\max\{0, I_0-N+n\}}^{\min\{n, I_0\}} \binom{I_0}{i} [q(t)]^i [1-q(t)]^{I_0-i} \times \binom{N-I_0}{n-i} [r(t)]^{n-i} [1-r(t)]^{N-I_0-(n-i)} \quad 1 \leq n \leq N-1, \quad (16)$$

$$P_N(t) = [q(t)]^{I_0} [r(t)]^{N-I_0} = \left[\frac{\nu + \mu e^{-(\nu+\mu)t}}{\nu + \mu} \right]^{I_0} \left[\frac{\nu (1 - e^{-(\nu+\mu)t})}{\nu + \mu} \right]^{N-I_0}. \quad (17)$$

We note that $P_0(t)$ is the probability that there are no infected persons and that $P_N(t)$ is the probability that all persons are infected at time t , respectively, for the first time.

4.3 Steady State

In the steady state at time $t \rightarrow \infty$, we have

$$q(\infty) \triangleq \lim_{t \rightarrow \infty} q(t) = \frac{\nu}{\nu + \mu} \quad ; \quad r(\infty) \triangleq \lim_{t \rightarrow \infty} r(t) = \frac{\nu}{\nu + \mu} = q(\infty). \quad (18)$$

Therefore, we get the following statistics.

$$G(z, \infty) \triangleq \lim_{t \rightarrow \infty} G(z, t) = \left(\frac{\nu z + \mu}{\nu + \mu} \right)^N \quad (\text{binomial distribution}). \quad (19)$$

$$P_n = \lim_{t \rightarrow \infty} P_n(t) = \binom{N}{n} \left(\frac{\nu}{\nu + \mu} \right)^n \left(\frac{\mu}{\nu + \mu} \right)^{N-n} \quad 0 \leq n \leq N. \quad (20)$$

$$E[I(\infty)] = \frac{N\nu}{\nu + \mu} \quad ; \quad \text{Var}[I(\infty)] = \frac{N\nu\mu}{(\nu + \mu)^2}. \quad (21)$$

In Table 2, we assume $\lambda = 0$. The numerical values have been obtained from the exact analytical solution given above.

5 Inversion of Laplace Transforms

We first try inversion of the Laplace transforms of the set of simultaneous differential equations for $\{P_n(t); 0 \leq n \leq N\}$ shown in eqs. (4)–(6), which are written as a set of algebraic equations in terms of Laplace transforms $\{P_n^*(s); 0 \leq n \leq N\}$ via

$$P_n^*(s) \triangleq \int_0^\infty e^{-st} P_n(t) dt \quad ; \quad \int_0^\infty e^{-st} \frac{dP_n(t)}{dt} dt = sP_n^*(s) - P_n(0) \quad 0 \leq n \leq N \quad (22)$$

as follows:

$$\frac{d\mathbf{P}(t)}{dt} = \mathbf{P}(t)\mathcal{Q} \quad ; \quad s\mathbf{P}^*(s) - \mathbf{P}(0) = \mathbf{P}^*(s)\mathcal{Q}, \quad (23)$$

Table 2: Probabilities $\{P_n(t); 0 \leq n \leq 8\}$, the mean $E[I(t)]$, and the variance $\text{Var}[I(t)]$. Parameters: $N = 8, I(0) = 5, \lambda = 0, \nu = 0.2, \mu = 0.1$.

t	$P_0(t)$	$P_1(t)$	$P_2(t)$	$P_3(t)$	$P_4(t)$	$P_5(t)$	$P_6(t)$	$P_7(t)$	$P_8(t)$	$E[I(t)]$	$\text{Var}[I(t)]$
0	0	0	0	0	0	1.000000	0	0	0	5.00000	0
1	0.000003	0.000146	0.003137	0.034146	0.190938	0.471277	0.248359	0.048711	0.003283	5.08639	0.82345
2	0.000026	0.000777	0.009367	0.058665	0.199849	0.351167	0.273425	0.098606	0.012047	5.15040	1.26983
3	0.000067	0.001487	0.013744	0.067947	0.192502	0.311302	0.272751	0.119631	0.020569	5.19781	1.51072
4	0.000105	0.001994	0.016073	0.070965	0.186091	0.293925	0.270874	0.133123	0.026851	5.23294	1.63989
5	0.000130	0.002287	0.017153	0.071635	0.181596	0.285142	0.270077	0.140935	0.031045	5.25896	1.70853
6	0.000145	0.002435	0.017580	0.071445	0.178493	0.280345	0.270042	0.145762	0.033752	5.27823	1.74453
7	0.000153	0.002498	0.017692	0.070979	0.176325	0.277600	0.270369	0.148894	0.035491	5.29251	1.76306
8	0.000156	0.002517	0.017664	0.070468	0.174789	0.275974	0.270818	0.150995	0.036618	5.30309	1.77231
9	0.000157	0.002516	0.017583	0.070003	0.173689	0.274985	0.271269	0.152439	0.037359	5.31093	1.77671
10	0.000157	0.002507	0.017489	0.069612	0.172893	0.274367	0.271669	0.153451	0.037855	5.31674	1.77863
∞	0.000152	0.000305	0.000610	0.001219	0.024390	0.004877	0.009755	0.019509	0.039018	5.33333	1.77778

which can be modified to

$$\mathbf{P}^*(s) (s\mathcal{I} - \mathcal{Q}) = \mathbf{P}(0) \quad ; \quad \mathbf{P}^*(s) = \mathbf{P}(0) (s\mathcal{I} - \mathcal{Q})^{-1}, \quad (24)$$

where $\mathbf{P}^*(s)$ is a row vector for $\{P_n^*(s); 0 \leq n \leq N\}$, $\mathbf{P}(0)$ is a row vector for the initial condition of a row vector $\mathbf{P}(t)$ for $\{P_n(t); 0 \leq n \leq N\}$ at $t = 0$, The **infinitesimal generator** \mathcal{Q} is an $(N + 1) \times (N + 1)$ tridiagonal (non-symmetric) matrix, of which the (m, n) th element is the state transition rate from state n to state m , $0 \leq m, n \leq N$. \mathcal{I} is the $(N + 1) \times (N + 1)$ identity matrix in which the diagonal elements are 1 and all off-diagonal elements are 0:

$$\mathbf{P}^*(s) \triangleq (P_0^*(s), P_1^*(s), \dots, P_n^*(s), \dots, P_N^*(s)), \quad (25)$$

$$\mathbf{P}(t) \triangleq (P_0(t), P_1(t), \dots, P_n(t), \dots, P_N(t)), \quad (26)$$

$$\mathcal{Q} \triangleq \begin{pmatrix} Q_{0,0} & Q_{0,1} & Q_{0,2} & \cdots & Q_{0,n} & \cdots & Q_{0,N-1} & Q_{0,N} \\ Q_{1,0} & Q_{1,1} & Q_{1,2} & \cdots & Q_{1,n} & \cdots & Q_{1,N-1} & Q_{1,N} \\ Q_{2,0} & Q_{2,1} & Q_{2,2} & \cdots & Q_{2,n} & \cdots & Q_{2,N-1} & Q_{2,N} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \vdots \\ Q_{N-1,0} & Q_{N-1,1} & Q_{N-1,2} & \cdots & Q_{N-1,n} & \cdots & Q_{N-1,N-1} & Q_{N-1,N} \\ Q_{N,0} & Q_{N,1} & Q_{N,2} & \cdots & Q_{N,n} & \cdots & Q_{N,N-1} & Q_{N,N} \end{pmatrix}, \quad (27)$$

where the nonzero elements of \mathcal{Q} are given by

$$Q_{n,n} = -(\lambda_n + \mu_n) = -\{n\lambda + (N - n)\nu + n\mu\}, \quad 1 \leq n \leq N - 1,$$

$$Q_{0,0} = -N\nu \quad ; \quad Q_{N,N} = -N\mu,$$

$$Q_{n,n+1} = \lambda_n \triangleq n\lambda + (N - n)\nu, \quad 0 \leq n \leq N - 1,$$

$$Q_{n,n-1} = \mu_n \triangleq n\mu, \quad 1 \leq n \leq N.$$

Since $Q_{n,n} + Q_{n,n+1} + Q_{n,n-1} = 0$, $1 \leq n \leq N-1$, and $Q_{0,0} + Q_{1,0} = Q_{N-1,N} + Q_{N,N} = 0$, the sum of all elements in every row (row sum) is zero.

The **stationary distribution** $\boldsymbol{\pi} = (\pi_0, \pi_1, \dots, \pi_N)$ (a row vector) can be obtained either from the detailed balance equations

$$\pi_n \lambda_n = \pi_{n+1} \mu_{n+1}, \quad 0 \leq n \leq N-1, \quad (28)$$

or as the left eigenvector of matrix \mathcal{Q} associated with eigenvalue 0, both along with the usual normalization:

$$\boldsymbol{\pi} \mathcal{Q} = 0\boldsymbol{\pi}, \quad \sum_{n=0}^N \pi_n = 1. \quad (29)$$

For our example with the same parameters as in Table 1, we have

$$\mathcal{Q} \triangleq \begin{pmatrix} -1.6 & 1.6 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0.1 & -2 & 1.9 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0.2 & -2.4 & 2.2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0.3 & -2.8 & 2.5 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.4 & -3.2 & 2.8 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0.5 & -3.6 & 3.1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0.6 & -4 & 3.4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0.7 & -4.4 & 3.7 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0.8 & -0.8 \end{pmatrix}, \quad (30)$$

Through the above-mentioned procedure, we obtain

$$\begin{aligned} P_0(t) &= 1.73732 \times 10^{-7} - 1.92016 \times 10^{-7} e^{s_1 t} - 7.6608 \times 10^{-6} e^{s_2 t} + 0.0000419876 e^{s_3 t} \\ &+ 0.0000216199 e^{s_4 t} - 0.000499537 e^{s_5 t} + 0.00117554 e^{s_6 t} - 0.00113797 e^{s_7 t} \\ &+ 0.000406046 e^{s_8 t}, \\ P_1(t) &= 2.77972 \times 10^{-6} + 9.23127 \times 10^{-6} e^{s_1 t} + 0.000254654 e^{s_2 t} - 0.000943395 e^{s_3 t} \\ &- 0.000304069 e^{s_4 t} + 0.0036748 e^{s_5 t} - 0.00237707 e^{s_6 t} - 0.00239809 e^{s_7 t} \\ &+ 0.00208116 e^{s_8 t}, \\ P_2(t) &= 0.0000264073 - 0.0002019 e^{s_1 t} - 0.0036619 e^{s_2 t} + 0.00837795 e^{s_3 t} + 0.00135717 e^{s_4 t} \\ &- 0.00217077 e^{s_5 t} - 0.0117552 e^{s_6 t} + 0.00178083 e^{s_7 t} + 0.00624636 e^{s_8 t}, \\ P_3(t) &= 0.000193654 + 0.00263863 e^{s_1 t} + 0.0291974 e^{s_2 t} - 0.0344438 e^{s_3 t} - 0.000817651 e^{s_4 t} \\ &- 0.0237394 e^{s_5 t} - 0.0083689 e^{s_6 t} + 0.0211877 e^{s_7 t} + 0.0141524 e^{s_8 t}, \\ P_4(t) &= 0.00121033 - 0.022687 e^{s_1 t} - 0.134906 e^{s_2 t} + 0.0441058 e^{s_3 t} - 0.00704244 e^{s_4 t} \\ &- 0.0156198 e^{s_5 t} + 0.0437773 e^{s_6 t} + 0.0649309 e^{s_7 t} + 0.0262311 e^{s_8 t}, \\ P_5(t) &= 0.00677788 + 0.132346 e^{s_1 t} + 0.319201 e^{s_2 t} + 0.115118 e^{s_3 t} + 0.00136186 e^{s_4 t} \\ &+ 0.0916948 e^{s_5 t} + 0.164228 e^{s_6 t} + 0.129206 e^{s_7 t} + 0.0400664 e^{s_8 t}, \\ P_6(t) &= 0.035019 - 0.513411 e^{s_1 t} - 0.0748694 e^{s_2 t} - 0.253279 e^{s_3 t} + 0.034212 e^{s_4 t} \\ &+ 0.266118 e^{s_5 t} + 0.287783 e^{s_6 t} + 0.173058 e^{s_7 t} + 0.0453692 e^{s_8 t}, \\ P_7(t) &= 0.170092 + 1.1797 e^{s_1 t} - 1.31476 e^{s_2 t} - 0.56505 e^{s_3 t} + 0.0425288 e^{s_4 t} \\ &+ 0.22666 e^{s_5 t} + 0.17626 e^{s_6 t} + 0.0732389 e^{s_7 t} + 0.0113342 e^{s_8 t}, \\ P_8(t) &= 0.786677 - 0.778394 e^{s_1 t} + 1.17956 e^{s_2 t} + 0.686073 e^{s_3 t} - 0.0713173 e^{s_4 t} \\ &- 0.546117 e^{s_5 t} - 0.650772 e^{s_6 t} - 0.459866 e^{s_7 t} - 0.145888 e^{s_8 t}, \end{aligned}$$

where $\{s_1 = -6.40755, s_2 = -4.92412, s_3 = -3.84732, s_4 = -3.00643, s_5 = -2.33564, s_6 = -1.80221, s_7 = -1.38927, s_8 = -1.08746\}$ are negative eigenvalues of matrix \mathcal{Q} . The stationary distribution is given by the left eigenvector: $\mathbf{u}_0 = \boldsymbol{\pi} = (1.73732 \times 10^{-7}, 2.77972 \times 10^{-6}, 0.0000264073, 0.000193654, 0.00121033, 0.00677788, 0.035019, 0.170092, 0.786677)$.

We can confirm that the expressions

$$\sum_{n=0}^8 P_n(t) = 1 \quad ; \quad E[I(t)] = \sum_{n=0}^8 n P_n(t) \quad ; \quad \text{Var}[I(t)] = \sum_{n=0}^8 (n - E[I(t)])^2 P_n(t) \quad (31)$$

yield the numerical values for each time t shown in Table 1.

6 Syski's Spectral Decomposition

Syski's spectral decomposition method is presented in his classic book [6, Chapter 5]. It can be applied to general continuous-time Markov processes. It was used by Kobayashi and Ren [5] for their transient analysis of a finite number of on-off sources in a statistical multiplexer of communication networks.

Let s_n be the n th eigenvalue of matrix \mathcal{Q} , and \mathbf{u}_n and \mathbf{v}_n be the left (row) and right (column) eigenvectors, respectively, associated with s_n :²

$$\mathbf{u}_n \mathcal{Q} = s_n \mathbf{u}_n, \quad \mathcal{Q} \mathbf{v}_n = s_n \mathbf{v}_n, \quad 0 \leq n \leq N. \quad (32)$$

The left eigenvectors $\{\mathbf{u}_n\}$ are calculated from the right eigenvectors $\{\mathbf{v}_n\}$ as

$$(\mathbf{u}_1^T, \mathbf{u}_2^T, \dots, \mathbf{u}_N^T) = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_N)^{-1} \quad (33)$$

so that they are **bi-orthogonal** and normalized as

$$\mathbf{u}_m \mathbf{v}_n = \sum_{k=0}^N u_m(k) v_n(k) = \delta_{m,n} = \begin{cases} 1 & m = n \\ 0 & m \neq n \end{cases}, \quad (34)$$

where $u_m(k)$ is the k th element of the m th left eigenvector \mathbf{u}_m and $v_n(k)$ is the k th element of the n th right eigenvector \mathbf{v}_n . We tabulate s_m , \mathbf{v}_m and \mathbf{u}_m in Tables 3 and 4.

In particular, since $s_0 = 0$, we have $\mathcal{Q} \mathbf{v}_0 = 0$ so that $\mathbf{v}_0 = (1, 1, \dots, 1)^T$, because all row sums are zero. On the other hand, since $\mathbf{u}_0 \mathcal{Q} = 0 \mathbf{u}_0$, we have the stationary distribution $\mathbf{u}_0 = \boldsymbol{\pi}$.

Now, we have the formal solution

$$\frac{d\mathbf{P}(t)}{dt} = \mathbf{P}(t) \mathcal{Q} \quad \implies \quad \mathbf{P}(t) = \mathbf{P}(0) e^{\mathcal{Q}t}, \quad (35)$$

where the **matrix exponential** is given by

$$e^{\mathcal{Q}t} = \sum_{n=0}^N e^{s_n t} \mathbf{v}_n \mathbf{u}_n = \mathbf{v}_0 \mathbf{u}_0 + \sum_{n=1}^N e^{s_n t} \mathbf{v}_n \mathbf{u}_n. \quad (36)$$

²In Mathematica, the right eigenvectors of a matrix Q are obtained by `Eigenvectors[Q]`, while the left eigenvectors are obtained by `Eigenvectors[Transpose[Q]]`.

If we define the following $n + 1$ scalars

$$\gamma_0 = 1 \quad ; \quad \gamma_n \triangleq \mathbf{P}(0)\mathbf{v}_n = \sum_{k=0}^N P_k(0)v_n(k), \quad 1 \leq n \leq N, \quad (37)$$

we get the **spectral decomposition** of $\mathbf{P}(t)$:

$$\begin{aligned} \mathbf{P}(t) &= \mathbf{P}(0)v_0\mathbf{u}_0 + \sum_{n=1}^N e^{s_n t} \mathbf{P}(0)v_n\mathbf{u}_n = \gamma_0\mathbf{u}_0 + \sum_{n=1}^N e^{s_n t} \gamma_n\mathbf{u}_n \\ &= \boldsymbol{\pi} + \sum_{n=1}^N e^{s_n t} \gamma_n\mathbf{u}_n, \quad t \geq 0. \end{aligned} \quad (38)$$

If the initial distribution vector $\mathbf{P}(0)$ is such that the only I_0 th element is 1, i.e., the process starts with state I_0 , the m th element of the row vector $\mathbf{P}(t)$ is given by

$$P_m(t) = \pi_m + \sum_{n=1}^N e^{s_n t} v_n(I_0)u_n(m), \quad 0 \leq m \leq N, \quad (39)$$

where $v_n(I_0)$ is the I_0 th element of the n th right eigenvector \mathbf{v}_n , and $u_n(m)$ is the m th element of the n th left eigenvector \mathbf{u}_n , respectively, associated with eigenvalue s_n . This calculation agrees with the results given in Section 5.

For example, in order to obtain the coefficient of $e^{s_1 t}$ in the expression for $P_2(t)$, we multiply $v_1(5) = 0.55162$ in Table 3 and $u_1(2) = -0.00036612$ in Table 4. The result is -0.0002019 , which agrees precisely with the corresponding term.

7 Karlin-McGregor's Symmetrization

For efficient application of Syski's method, we refer to the Karlin-McGregor's symmetrization for birth-and-death processes, which was originally derived in [3] for general processes and later expressed in a matrix formulation by Keilson [4, Chapter 2] for processes with a finite number of states. The symmetrization follows from the reversibility of the birth-and-death process and the associated detailed balance equations in (28). In terms of the generator matrix \mathcal{Q} , the detailed balance equations can be written as

$$\pi_n \mathcal{Q}_{n,n+1} = \mathcal{Q}_{n+1,n} \pi_{n+1}, \quad 0 \leq n \leq N-1. \quad (40)$$

We define the diagonal matrix $D \triangleq \text{diag}(\pi_0, \pi_1, \dots, \pi_N)$ in terms of the stationary distribution. Then, eq. (28) implies that

$$D\mathcal{Q} = \mathcal{Q}^T D = (D\mathcal{Q})^T, \quad (41)$$

i.e., the matrix $D\mathcal{Q}$ is symmetric. We further define the matrix

$$J \triangleq D^{1/2} \mathcal{Q} D^{-1/2}, \quad (42)$$

Table 3: Eigenvalues $\{s_m\}$ and the n th element of the m th right eigenvector $v_m(n)$ of \mathcal{Q} . All the elements of \mathbf{v}_0 are equal (set to 1) because each row sum of matrix \mathcal{Q} is zero.

m	s_m	$v_m(0)$	$v_m(1)$	$v_m(2)$	$v_m(3)$	$v_m(4)$	$v_m(5)$	$v_m(6)$	$v_m(7)$	$v_m(8)$
0	0	1	1	1	1	1	1	1	1	1
1	-6.40755	-0.031223	0.093817	-0.215991	0.384924	-0.529534	0.551622	-0.414174	0.195934	-0.027953
2	-4.92412	-0.170347	0.353908	-0.535702	0.582452	-0.430594	0.181933	-0.008259	-0.029861	0.005792
3	-3.84732	-0.435713	0.611992	-0.572092	0.320729	-0.065712	-0.030627	0.013042	0.005990	-0.001573
4	-3.00643	-0.716733	0.630022	-0.296000	0.024318	-0.033512	0.001157	-0.005627	-0.001440	0.000522
5	-2.33564	-0.907563	0.417275	-0.025946	-0.038693	-0.004073	-0.004270	0.002399	0.000421	-0.000219
6	-1.80221	-0.989961	0.125113	0.065127	0.006322	-0.005292	-0.003545	-0.001202	-0.000152	0.000121
7	-1.38927	-0.991213	-0.130551	0.010205	0.016557	0.008118	0.002885	0.000748	0.000065	-0.000088
8	-1.08746	0.947481	0.303515	0.095907	0.029627	0.008786	0.002396	0.000525	0.000027	-0.000075

Table 4: The n th element of the m th left eigenvector $u_m(n)$ of \mathcal{Q} , where $u_0(n) = \pi_n$ (stationary distribution).

m	$u_m(0)$	$u_m(1)$	$u_m(2)$	$u_m(3)$	$u_m(4)$	$u_m(5)$	$u_m(6)$	$u_m(7)$	$u_m(8)$
0	1.74×10^{-7}	2.78×10^{-6}	0.000026	0.000194	0.001210	0.006778	0.035019	0.170092	0.786677
1	3.5×10^{-7}	1.67×10^{-5}	-0.00037	0.004783	-0.04113	0.239923	-0.930729	2.138603	-1.411102
2	-4.2×10^{-5}	0.0014	-0.02013	0.160484	-0.74152	1.754495	-0.411522	-7.22663	6.483459
3	-0.00137	0.030803	-0.27355	1.12463	-1.44011	-3.75873	8.26986	18.4496	-22.4011
4	-0.01868	0.262755	-1.17276	0.706554	6.08556	-1.176820	-29.563488	-36.7503	61.627201
5	-0.11698	0.860583	-0.050836	-5.55942	-3.65793	21.473559	62.320822	53.08028	-127.8925
6	-0.33161	0.670549	3.316013	2.360783	-12.3492	-46.326962	-81.180816	-49.7211	183.56229
7	-0.39448	-0.8313	0.617329	7.344756	22.50841	44.789676	59.990771	25.38842	-159.4136
8	0.169439	0.868447	2.606961	5.905663	10.94596	16.719304	18.932117	4.729649	-60.87754

where

$$D^{1/2} \triangleq \text{diag}(\sqrt{\pi_0}, \sqrt{\pi_1}, \dots, \sqrt{\pi_N}) \quad ; \quad D^{-1/2} \triangleq \text{diag}\left(\frac{1}{\sqrt{\pi_0}}, \frac{1}{\sqrt{\pi_1}}, \dots, \frac{1}{\sqrt{\pi_N}}\right). \quad (43)$$

We can verify that J is symmetric using eq. (40) as follows:

$$J^T = D^{-1/2} \mathcal{Q}^T D^{1/2} = D^{-1/2} (\mathcal{Q}^T D) D^{-1/2} = D^{-1/2} (D \mathcal{Q}) D^{-1/2} = D^{1/2} \mathcal{Q} D^{-1/2} = J. \quad (44)$$

Since J is symmetric, it has real eigenvalues and an orthonormal basis of (right) eigenvectors $\{\mathbf{w}_0, \mathbf{w}_1, \dots, \mathbf{w}_N\}$. The left eigenvectors are simply $\{\mathbf{w}_0^T, \mathbf{w}_1^T, \dots, \mathbf{w}_N^T\}$.

The matrices J and \mathcal{Q} have the same diagonal elements and J retains the tridiagonal form of \mathcal{Q} . Noting that

$$\mathcal{Q} \mathbf{v}_n = s_n \mathbf{v}_n \quad \implies \quad D^{1/2} \mathcal{Q} \mathbf{v}_n = s_n D^{1/2} \mathbf{v}_n, \quad (45)$$

and letting $\mathbf{w}_n \triangleq D^{1/2} \mathbf{v}_n$, we obtain

$$D^{1/2} \mathcal{Q} D^{-1/2} \mathbf{w}_n = s_n \mathbf{w}_n \quad \implies \quad J \mathbf{w}_n = s_n \mathbf{w}_n, \quad 0 \leq n \leq N. \quad (46)$$

Thus, J has the same eigenvalues $\{s_0, s_1, s_2, \dots, s_N\}$ as Q . The right and left eigenvectors of Q can be obtained from the eigenvalues of J as follows:

$$\mathbf{v}_n = D^{-1/2}\mathbf{w}_n, \quad ; \quad \mathbf{u}_n = \mathbf{w}_n^T D^{1/2}, \quad 0 \leq n \leq N, \quad (47)$$

respectively. Since J is symmetric, numerical computation of the eigenvectors of J is more efficient and stable than computation of the eigenvectors of Q [2, Chapter 8, pp.408–412]. Therefore, for efficient numerical computation of $\mathbf{P}(t)$ in Syski’s method, we apply the symmetrization transformation given by (41) and then compute the decomposition of J to obtain the eigenvalues $\{s_n\}$ and the eigenvectors $\{\mathbf{w}_n\}$. The spectral decomposition in (37) can then be applied using the relations in (47).

References

- [1] N. T. J. Bailey, *The Elements of Stochastic Processes with Applications to Natural Sciences*, John Wiley & Sons, Inc., 1964.
- [2] G. H. Golub and C. F. Van Loan, *Matrix Computations*, Fourth edition, The Johns Hopkins University Press, Baltimore, 2013.
- [3] S. Karlin and J. L. McGregor, Coincidence properties of birth and death processes, *Pacific Journal of Mathematics*, Vol.9, No.4, pp.1109–1140, August 1959.
- [4] J. Keilson, *Markov Chain Models — Rarity and Exponentiality*, Springer Lecture Notes in Mathematics, Vol.28, Springer, 2013; First edition, 1979.
- [5] H. Kobayashi and Q. Ren, A mathematical theory for transient analysis of communication networks (invited paper), *IEICE Transactions on Communications*, vol. E-75B, no. 12, pp. 1266–1276, December 1992.
- [6] R. Syski, *Introduction to Congestion Theory in Telephone Systems*, Second edition, North-Holland, Amsterdam, 1986; First edition, Oliver and Boyd, Edinburgh, 1960.
- [7] T. Yamagishi, H. Kamiya, K. Kakimoto, M. Suzuki, and T. Wakita, Descriptive study of COVID-19 outbreak among passengers and crew on Diamond Princess cruise ship, Yokohama Port, Japan, 20 January to 9 February 2020, *Eurosurveillance* 25(23), 2000272, 2020 June, <https://pubmed.ncbi.nlm.nih.gov/32191689>

Acknowledgments

This paper is dedicated to the memory of the late Hisashi Kobayashi, the Sherman Fairchild University Professor of Electrical Engineering and Computer Science, Emeritus at Princeton University, who passed away on March 9, 2023 at the age of 84.

The authors used ChatGPT (Open AI) as an assistive tool for drafting the text. However, all results developed and verified are the responsibility of the authors.

This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.